Neuroscience Letters, 277–280
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Akaike, N., see Oyama, Ya., 153
Alberch, J., Brito, B., Notario, V. and Castro, R., Prenatal haloperidol treatment decreases nerve growth factor receptor and mRNA in neonate rat forebrain, 228
Alexander, S.P.H., Hill, S.J. and Kendall, D.A., Do polyamines regulate the NMDA inhibition of muscarinic receptor-induced phosphoinositide turnover in guinea pig brain?, 167
Allaoua, H., see Chaudieu, I., 263
Allen, S.J., MacGowan, S.H., Treanor, J.J.S., Feeney, R., Wilcock, G.K. and Dawbarn, D., Normal β-NGF content in Alzheimer's disease cerebral cortex and hippocampus, 135
Alvarez, A., see Verney, C., 121
Andou, Ya., see Mitani, A., 171
Anwyl, R., see McGuinness, N., 13
Arai, Ya., see Matsumoto, A., 159
Ardelt, B.K., see Patel, M.N., 42
Arvidsson, U., see Meister, B., 66
Azuma, H., Yamatodani, A., Yagi, A., Nishimura, T. and Wada, H., Effect of bright light in the morning on diurnal variations of pineal indoles in NZBWF1 mice, 210

Bagetta, G., see Constanti, A., 27
Bahr, B.A., Vanderklish, P.W., Ha, L.T., Tin, M.T. and Lynch, G., Spectrin breakdown products increase with age in telencephalon of mouse brain, 237
Barret, L., Torch, S., Usson, Y., Gonthier, B. and Saxod, R., A morphometric evaluation of the effects of trichloroethylene and dichloroacetylene on the rat mental nerve. Preliminary results, 141
Beal, J.A., see Nandi, K.N., 8
Bel, E.A.D. and Slater, P., Binding to the glycine site of the NMDA receptor complex in brains of patients with Alzheimer's disease, 75
Bellaidi, M., see Chaudieu, I., 263
Berger, B., see Verney, C., 121
Bernardi, G., see Mercuri, N.B., 145
Bigot, D. and Hunt, S.P., The effects of quisqualate and nocodazole on the organization of MAP2 and neurofilaments in spinal cord neurons in vitro, 21
Boulu, R.G., see Buisson, A., 257
Brito, B., see Alberch, J., 228
Brotchie, P., Iansek, R. and Horne, M., A neural network model of neural activityin the monkey globus pallidus, 33
Brundell, P., Jr., R.G., Kerry, C.J., Nakanishi, K., Sudan, H.L. and Usherwood, P.N.R., Quisqualate-sensitive glutamate receptors of the locust *Schistocerca gregaria* are antagonised by intracellularly applied philanthotoxin and spermine, 196
Buisson, A., Pateau, V., Plotkine, M. and Boulu, R.G., Nigrostriatal pathway modulates striatum vulnerability to quinolinic acid, 257

Calabresi, P., see Mercuri, N.B., 145
Cameron, A.A., Cliffer, K.D., Dougherty, P.M., Willis, W.D. and Carlton, S.M., Changes in lectin, GAP-43 and neuropeptide staining in the rat superficial dorsal horn following experimental peripheral neuropathy, 249
Carlton, S.M., Dougherty, P.M., Pover, C.M. and Coggeshall, R.E., Neuroma formation and numbers of axons in a rat model of experimental peripheral neuropathy, 88
Carlton, S.M., see Cameron, A.A., 249
Carson, J., see Westlund, K.N., 183
Casanova, M.F., see Laruelle, M., 273
Castro, R., see Alberch, J., 228
Chaudieu, I., Allaoua, H., Bellaidi, M., Rondouin, G., Kamenka, J.M. and Chicheportiche, R., Effect of amygdaloid kindling on the high- and low-affinity [^{3}H]TCP binding sites of the rat CNS, 263
Chicheportiche, R., see Chaudieu, I., 263
Clerin, E., see Evercooren, A.B.-V., 241
Cliffer, K.D., see Cameron, A.A., 249
Coderre, T.J. and Melzack, R., Central neural mediators of secondary hyperalgesia following heat injury in rats: neuropeptides and excitatory amino acids, 71
Coggeshall, R.E., Reynolds, M.L. and Woolf, C.J., Distribution of the growth associated protein GAP-43 in the central processes of axotomized primary afferents in the adult rat spinal cord; presence of growth cone-like structures, 37
Coggeshall, R.E., see Carlton, S.M., 88
Constanti, A. and Bagetta, G., Muscarinic receptor activation induces a prolonged post-stimulus afterdepolarization with a conductance decrease in guinea-pig olfactory cortex neurones in vitro, 27
Cordell, B., see Jr., G.M.M., 100
Couraud, P.O., see Durieu-Trautmann, O., 175
Crossman, A.R., see Graham, W.C., 129
Csillik, B., see Knyihár-Csillik, E., 83
Cubeddu, L.X., Hoffmann, I.S., Talmaciu, R.K., Diliberto, P. and Lovemberg, T., Carbachol–phorbol ester interactions at muscarinic receptors modulating dopamine release from rabbit striatal slices, 245

Dawbarn, D., see Allen, S.J., 135
Deguchi, T., see Ichikawa, T., 213
Diliberto, P., see Cubeddu, L.X., 245
DiPette, D.J., see Westlund, K.N., 183
Dockray, G.J., see Forster, E.R., 5
Dougherty, P.M., see Cameron, A.A., 249
Dougherty, P.M., see Carlton, S.M., 88
Drescher, D.G., see Khan, K.M., 109
Drescher, M.J., see Khan, K.M., 109
Dulhunty, A.F., Junankar, P.R. and Stanhope, C., Calcium ATPase in the sarcoplasmic reticulum of muscle from normal and malignant hyperthermia susceptible pigs, 187
Durieu-Trautmann, O., Couraud, P.O., Foignant-Chaverot, N. and Strosberg, A.D., cAMP-dependent down-regulation of endothelin-1 receptors on rat astrocytoma C6 cells, 175

Ellis, W.G., see Jr., G.M.M., 100
Eng, L.F., see Jr., G.M.M., 100
Evercooren, A.B.-V., Gansmüller, A., Clerin, E. and Gumpel, M., Hoechst 33342 a suitable fluorescent marker for Schwann cells after transplantation in the mouse spinal cord, 241
Eysel, U.T. and Schmidt-Kastner, R., Neuronal dysfunction at the border of focal lesions in cat visual cortex, 45

Farooqui, A.A., Wallace, L.J. and Horrõcks, L.A., Stimulation of mono- and diacylglycerol lipase activities in ibotenate-induced lesions of nucleus basalis magnocellularis, 97
Feeney, R., see Allen, S.J., 135

Foignant-Chaverot, N., see Durieu-Trautmann, O., 175
Formenti, A. and Sansone, V., Inhibitory action of acetylcholine, baclofen and GTP-γ-*S* on calcium channels in adult rat sensory neurons, 267
Forno, L.S., see Jr., G.M.M., 100
Forster, E.R. and Dockray, G.J., 5

Gansmüller, A., see Evercooren, A.B.-V., 241
Gebhart, G.F., see Ness, T.J., 113
Gillard, J., see Knight, A.R., 233
Gonthier, B., see Barret, L., 141
Gonzalez-DeWhitt, P.A., see Jr., G.M.M., 100
Graham, W.C., Hill, D.R., Hill,, , Woodruff, G.N., Sambrook, M.A. and Crossman, A.R., Reduction of [^{125}I]Bolton Hunter CCK8 and [^{3}H]MK-329 (devazepide) binding to CCK receptors in the substantia nigra/VTA complex and its forebrain projection areas following MPTP-induced hemi-parkinsonism in the monkey, 129
Greenberg, B.D., see Jr., G.M.M., 100
Greengard, P., see Meister, B., 66
Greenwood, C.E., see Tatton, W.G., 179
Gribkoff, V.K., Electrophysiological evidence for *N*-methyl-D-aspartate excitatory amino acid receptors in the rat supraoptic nucleus in vitro, 260
Guillamon, A., see Luque, J.M., 163
Gumpel, M., see Evercooren, A.B.-V., 241

Ha, L.T., see Bahr, B.A., 237
Hara, N., see Sawada, M., 217
Harata, N., see Oyama, Ya., 153
Hatfield, J.S., see Khan, K.M., 109
Henry, S. and Tappaz, M., Relative enrichment of the lighter 59 kDa form of glutamic acid decarboxylase in nerve endings: an immunoblotting study in pituitary neurointermediate lobe, 253
Higashi, S., see Murakami, F., 49
Hill,, , see Graham, W.C., 129
Hill, D.R., see Graham, W.C., 129
Hill, S.J., see Alexander, S.P.H., 167
Hoffmann, I.S., see Cubeddu, L.X., 245
Hökfelt, T., see Meister, B., 66
Holland, O.B., see Westlund, K.N., 183
Horne, M., see Brotchie, P., 33
Horrocks, L.A., see Farooqui, A.A., 97
Hösli, E., see Hösli, L., 193
Hösli, L., Hösli, E., Lefkovits, M. and Wagner, S., Electrophysiological evidence for the existence of receptors for endothelin and vasopressin on cultured astrocytes of rat spinal cord and brainstem, 193
Huang, L.-Y.M., see McPherson, D.R., 201
Hunt, S.P., see Bigot, D., 21
Hwang, B.H., see Luque, J.M., 163
Hyodo, S., see Matsumoto, A., 159

Iansek, R., see Brotchie, P., 33
Ichikawa, T., Ohsako, S. and Deguchi, T., Production of an antiserum using a fusion protein produced by a cDNA for rat choline acetyltransferase, 213
Ikai, Y., see Shinonaga, Ya., 53
Immunoneutralization suggests that calcitonin gene related peptide regulates gastric emptying in the rat, 5
Ishizaki, Ya. and Murota, S., Arachidonic acid metabolism in cultured astrocytes: presence of 12-lipoxygenase activity in the intact cells, 149
Isom, G.E., see Patel, M.N., 42
Ito, S., see Wakamatsu, K., 57

Itoh, K., see Shinonaga, Ya., 53

Jacobowitz, D.M., see Yamaguchi, T., 61
Jacobowitz, D.M., see Yamaguchi, T., 79
Jr., G.M.M., Murphy, E., Greenberg, B.D., Cordell, B., Eng, L.F., Ellis, W.G., Forno, L.S., Salamat, S.M., Gonzalez-DeWhitt, P.A., Lowery, D.E. and Tinklenberg, J.R., Alzheimer's disease: β-amyloid precursor protein expression in plaques varies among cytoarchitectonic areas of the medial temporal lobe, 100
Jr., H.C.H., see Meister, B., 66
Jr., R.G., see Brundell, P., 196
Junankar, P.R., see Dulhunty, A.F., 187

Kamenka, J.M., see Chaudieu, I., 263
Kamo, I., see Kobayashi, T., 221
Kaneko, T., see Shinonaga, Ya., 53
Kataoka, K., see Mitani, A., 171
Kendall, D.A., see Alexander, S.P.H., 167
Kerry, C.J., see Brundell, P., 196
Khan, K.M., Hatfield, J.S., Drescher, M.J. and Drescher, D.G., The histochemical localization of acetylcholinesterase in the rainbow trout saccular macula by electron microscopy, 109
Kleinman, J.E., see Laruelle, M., 273
Knight, A.R., Gillard, J., Wong, E.H.F. and Middlemiss, D.N., The human σ site, which resembles that in NCB20 cells, may correspond to a low-affinity site in guinea pig brain, 233
Knight, D.S., see Nandi, K.N., 8
Knyihár-Csillik, E., Kreutzberg, G.W., Raivich, G. and Csillik, B., Vasoactive intestinal polypeptide in dorsal root terminals of the rat spinal cord is regulated by the axoplasmic transport in the peripheral nerve, 83
Kobayashi, T., Sano, M., Tsukagoshi, H. and Kamo, I., Muscle inactivity reduces the content of parvalbumin in rat thymic myoid cells in vitro, 221
Kreutzberg, G.W., see Knyihár-Csillik, E., 83

Laruelle, M., Sidhu, A., Casanova, M.F., Weinberger, D.R. and Kleinman, J.E., Characterization of [^{125}I]SCH 23982 binding in human brain: comparison with [^{3}H]SCH 23390, 273
Lefkovits, M., see Hösli, L., 193
Lethco, B.A., see Suchy, S.F., 105
Li, Yu.Q., see Zeng, S.-L., 205
Lo, E.H., Sun, G.-H. and Steinberg, G.K., Effects of NMDA and calcium channel antagonists on regional cerebral blood flow, 17
Lovemberg, T., see Cubeddu, L.X., 245
Lowery, D.E., see Jr., G.M.M., 100
Luque, J.M., Guillamon, A. and Hwang, B.H., Quantitative autoradiographic study on tyrosine hydroxylase mRNA with in situ hybridization and α_2 adrenergic receptor binding in the locus coeruleus of the spontaneously hypertensive rat, 163
Lynch, G., see Bahr, B.A., 237

MacGowan, S.H., see Allen, S.J., 135
Maeno, T., see Sawada, M., 217
Masuda, S., see Mitani, A., 171
Matsumoto, A., Arai, Ya., Urano, A. and Hyodo, S., Androgen regulates gap junction mRNA expression in androgen-sensitive motoneurons in the rat spinal cord, 159
McGuinness, N., Anwyl, R. and Rowan, M.,
McPherson, D.R. and Huang, L.-Y.M., Voltage-dependent tetrodotoxin binding to single batrachotoxin-modified Na channels recorded from intact neuroblastoma cells, 201
Meister, B., Arvidsson, U., Jr., H.C.H., Greengard, P. and Hökfelt,

T., Dopamine- and adenosine-3′:5′-monophosphate(cAMP)-regulated phosphoprotein of M_r 32,000 (DARPP-32) in the retina of cat, monkey and human, 66
Melzack, R., see Coderre, T.J., 71
Mercuri, N.B., Stratta, F., Calabresi, P. and Bernardi, G., Nomifensine but not amantadine increases dopamine-induced responses on rat substantia nigra zona compacta neurons, 145
Middlemiss, D.N., see Knight, A.R., 233
Mitani, A., Andou, Ya., Masuda, S. and Kataoka, K., Transient forebrain ischemia of three-minute duration consistently induces severe neuronal damage in field CA1 of the hippocampus in the normothermic gerbil, 171
Mizuno, N., see Shinonaga, Ya., 53
Murakami, F., Saito, Y., Higashi, S. and Oikawa, H., Synapses formed by ectopic corticofugal axons: an electron microscopic study of crossed corticorubral projections in kittens, 49
Murota, S., see Ishizaki, Ya., 149
Murphy, E., see Jr., G.M.M., 100

Nagai, T. and Oka, Y., The glossopharyngeal nerve of the axolotl labeled with carbocyanine dye (diI), 125
Nagatsu, T., see Wakamatsu, K., 57
Nakanishi, K., see Brundell, P., 196
Nandi, K.N., Knight, D.S. and Beal, J.A., Neurogenesis of ascending supraspinal projection neurons:ipsi versus contralateral projections, 8
Ness, T.J., Randich, A. and Gebhart, G.F., Further behavioral evidence that colorectal distension is a 'noxious' visceral stimulus in rats, 113
Nikolic, B., see Verney, C., 121
Nishimura, T., see Azuma, H., 210
Notario, V., see Alberch, J., 228

Ohsako, S., see Ichikawa, T., 213
Oikawa, H., see Murakami, F., 49
Oka, Y., see Nagai, T., 125
Oyama, Ya., Harata, N. and Akaike, N., Rapid interaction of tetraethylammonium with delayed K channel in mammalian brain neuron, 153

Pateau, V., see Buisson, A., 257
Patel, M.N., Ardelt, B.K., Yim, G.K.W. and Isom, G.E., Cyanide induces Ca^2-dependent and -independent release of glutamate from mouse brain slices, 42
Plotkine, M., see Buisson, A., 257
Pover, C.M., see Carlton, S.M., 88

Raivich, G., see Knyihár-Csillik, E., 83
Ramirez, J.J., see Suchy, S.F., 105
Randich, A., see Ness, T.J., 113
Rao, Z.-R., see Zeng, S.-L., 205
Reynolds, M.L., see Coggeshall, R.E., 37
Rondouin, G., see Chaudieu, I., 263
Rowan, M., see McGuinness, N., 13
Rusakov, D.A., Skibo, G.G. and Vasilenko, D.A., The spatial pattern of the synaptic vesicular apparatus as a correlate of transmitter storage models, 156

Saito, Y., see Murakami, F., 49
Salamat, S.M., see Jr., G.M.M., 100
Salo, P.T., see Tatton, W.G., 179
Sambrook, M.A., see Graham, W.C., 129
Sanides-Kohlrausch, C. and Wahle, P., Distribution and morphology of substance P-immunoreactive structures in the olfactory bulb and olfactory peduncle of the common marmoset (*Callithrix jacchus*), a primate species, 117
Sano, M., see Kobayashi, T., 221
Sansone, V., see Formenti, A., 267
Sawada, M., Hara, N. and Maeno, T., Tumor necrosis factor reduces the ACh-induced outward current in identified *Aplysia* neurons, 217
Saxod, R., see Barret, L., 141
Schmidt-Kastner, R., see Eysel, U.T., 45
Schwarting, G.A., see Suchy, S.F., 105
Seniuk, N.A., see Tatton, W.G., 179
Shi, J.-W., see Zeng, S.-L., 205
Shimizu, H., Alteration in hypothalamic monoamine metabolism of freely moving diabetic rat, 225
Shinonaga, Ya., Itoh, K., Takada, M., Kaneko, T., Ikai, Y. and Mizuno, N., Cholinergic projections from the globus pallidus to the lateral amygdaloid nucleus in the cat, 53
Sidhu, A., see Laruelle, M., 273
Skibo, G.G., see Rusakov, D.A., 156
Slater, P., see Bel, E.A.D., 75
Stanhope, C., see Dulhunty, A.F., 187
Steinberg, G.K., see Lo, E.H., 17
Stratta, F., see Mercuri, N.B., 145
Strosberg, A.D., see Durieu-Trautmann, O., 175
Suchy, S.F., Schwarting, G.A., Lethco, B.A. and Ramirez, J.J., The expression of a fucosyl-ganglioside in the molecular layer of the dentate gyrus following entorhinal cortical lesions, 105
Sudan, H.L., see Brundell, P., 196
Sun, G.-H., see Lo, E.H., 17

Takada, M., see Shinonaga, Ya., 53
Talmaciu, R.K., see Cubeddu, L.X., 245
Tappaz, M., see Henry, S., 253
Tatton, W.G., Greenwood, C.E., Salo, P.T. and Seniuk, N.A., Transmitter synthesis increases in substantia nigra neurons of the aged mouse, 179
The effects of external calcium on the *N*-methyl-D-aspartate induced short-term potentiation in the rat hippocampal slice, 13
Timmerman, W. and Westerink, B.H.C., Importance of the calcium content infused during microdialysis for the effects induced by D_2 agonists on the release of dopamine in the striatum of the rat, 93
Tinklenberg, J.R., see Jr., G.M.M., 100
Tin, M.T., see Bahr, B.A., 237
Torch, S., see Barret, L., 141
Treanor, J.J.S., see Allen, S.J., 135
Tsukagoshi, H., see Kobayashi, T., 221

Urano, A., see Matsumoto, A., 159
Usherwood, P.N.R., see Brundell, P., 196
Usson, Y., see Barret, L., 141

Vanderklish, P.W., see Bahr, B.A., 237
Vasilenko, D.A., see Rusakov, D.A., 156
Verney, C., Zecevic, N., Nikolic, B., Alvarez, A. and Berger, B., Early evidence of catecholaminergic cell groups in 5- and 6-week-old human embryos using tyrosine hydroxylase and dopamine-β-hydroxylase immunocytochemistry, 121

Wada, H., see Azuma, H., 210
Wagner, S., see Hösli, L., 193
Wahle, P., see Sanides-Kohlrausch, C., 117
Wakamatsu, K., Ito, S. and Nagatsu, T., Cysteinyldopamine is not incorporated into neuromelanin, 57

Wallace, L.J., see Farooqui, A.A., 97
Weinberger, D.R., see Laruelle, M., 273
Westerink, B.H.C., see Timmerman, W., 93
Westlund, K.N., DiPette, D.J., Carson, J. and Holland, O.B., Decreased spinal cord content of calcitonin gene-related peptide in the spontaneously hypertensive rat, 183
Wilcock, G.K., see Allen, S.J., 135
Willis, W.D., see Cameron, A.A., 249
Winsky, L., see Yamaguchi, T., 61
Winsky, L., see Yamaguchi, T., 79
Wong, E.H.F., see Knight, A.R., 233
Woodruff, G.N., see Graham, W.C., 129
Woolf, C.J., see Coggeshall, R.E., 37
Yagi, A., see Azuma, H., 210
Yamaguchi, T., Winsky, L. and Jacobowitz, D.M., Calretinin, a neuronal calcium binding protein inhibits phosphorylation of a 39 kDa synaptic membrane protein from rat brain cerebral cortex, 61
Yamaguchi, T., Winsky, L. and Jacobowitz, D.M., Calretinin, a neuronal calcium binding protein inhibits phosphorylation of a 39 kDa synaptic membrane protein from rat brain cerebral cortex, 79
Yamatodani, A., see Azuma, H., 210
Yim, G.K.W., see Patel, M.N., 42
Zecevic, N., see Verney, C., 121
Zeng, S.-L., Li, Yu.Q., Rao, Z.-R. and Shi, J.-W., Projections from serotonin- and substance P-like immunoreactive neurons in the midbrain periaqueductal gray onto the nucleus reticularis gigantocellularis pars α in the rat, 205

ANNOUNCEMENT

The Satellite Symposium of the IXth International Congress of Endocrinology 'Melatonin and the Pineal Gland: From Basic Science to Clinical Application' will be held in Paris, France, from 6 to 9 September 1991.

For further information:

Pr Y. Touitou
Department of Biochemistry
Faculty of Medicine Pitié-Salpetrière
91 boulevard de l'Hopital
75013 Paris
France

CONTENTS *(continued from back cover)*

The human σ site, which resembles that in NCB20 cells, may correspond to a low-affinity site in guinea pig brain
A.R. Knight, J. Gillard, E.H.F. Wong and D.N. Middlemiss (U.K.) 233
Spectrin breakdown products increase with age in telencephalon of mouse brain
B.A. Bahr, P.W. Vanderklish, L.T. Ha, M.T. Tin and G. Lynch (U.S.A.) 237
Hoechst 33342 a suitable fluorescent marker for Schwann cells after transplantation in the mouse spinal cord
A.B.-V. Evercooren, A. Gansmüller, E. Clerin and M. Gumpel (France) 241
Carbachol–phorbol ester interactions at muscarinic receptors modulating dopamine release from rabbit striatal slices
L.X. Cubeddu, I.S. Hoffmann, R.K. Talmaciu, P. Diliberto and T. Lovemberg (Venezuela, U.S.A.) 245
Changes in lectin, GAP-43 and neuropeptide staining in the rat superficial dorsal horn following experimental peripheral neuropathy
A.A. Cameron, K.D. Cliffer, P.M. Dougherty, W.D. Willis and S.M. Carlton (U.S.A.) 249
Relative enrichment of the lighter 59 kDa form of glutamic acid decarboxylase in nerve endings: an immunoblotting study in pituitary neurointermediate lobe
S. Henry and M. Tappaz (France) 253
Nigrostriatal pathway modulates striatum vulnerability to quinolinic acid
A. Buisson, V. Pateau, M. Plotkine and R.G. Boulu (France) 257
Electrophysiological evidence for *N*-methyl-D-aspartate excitatory amino acid receptors in the rat supraoptic nucleus in vitro
V.K. Gribkoff (U.S.A.) 260
Effect of amygdaloid kindling on the high- and low-affinity [^{3}H]TCP binding sites of the rat CNS
I. Chaudieu, H. Allaoua, M. Bellaidi, G. Rondouin, J.M. Kamenka and R. Chicheportiche (France) 263
Inhibitory action of acetylcholine, baclofen and GTP-γ-*S* on calcium channels in adult rat sensory neurons
A. Formenti and V. Sansone (Italy) 267
Characterization of [^{125}I]SCH 23982 binding in human brain: comparison with [^{3}H]SCH 23390
M. Laruelle, A. Sidhu, M.F. Casanova, D.R. Weinberger and J.E. Kleinman (U.S.A.) 273

Author index 277